

Title:

Persistent organic pollutant burden, experimental POP exposure and tissue properties affect metabolic profiles of blubber from grey seal pups.

Authors:

Kelly J. Robinson^{1*}, Ailsa J. Hall¹, Cathy Debier², Gauthier Eppe³, Jean-Pierre Thomé⁴ and Kimberley A. Bennett⁵

Affiliations:

¹Sea Mammal Research Unit, Scottish Oceans Institute, University of St Andrews

²Louvain Institute of Biomolecular Science and Technology, Université Catholique de Louvain

³Center for Analytical Research and Technology (CART), B6c, Department of Chemistry, Université de Liège

⁴Center for Analytical Research and Technology (CART), Laboratory of Animal Ecology and Ecotoxicology (LEAE), Université de Liège

⁵Division of Science, School of Science Engineering and Technology, Abertay University

*Corresponding Author: Kelly J. Robinson, email: kjr33@st-andrews.ac.uk


Keywords:

Dioxin-like PCBs; glucose uptake; lipolysis; lactate production; blubber depth; energetic state; environmental contamination; fasting; feeding

Abstract

Persistent organic pollutants (POPs) are toxic, ubiquitous, resist breakdown, bioaccumulate in living tissue and biomagnify in food webs. POPs can also alter energy balance in humans and wildlife. Marine mammals experience high POP concentrations, but consequences for their tissue metabolic characteristics are unknown. We used blubber explants from wild, grey seal (*Halichoerus grypus*) pups to examine impacts of intrinsic tissue POP burden and acute experimental POP exposure on adipose metabolic characteristics. Glucose use, lactate production and lipolytic rate differed between matched inner and outer blubber explants from the same individuals and between feeding and natural fasting. Glucose use decreased with blubber dioxin-like PCBs (DL-PCB) and increased with acute experimental POP exposure. Lactate production increased with DL-PCBs during feeding, but decreased with DL-PCBs during fasting. Lipolytic rate increased with blubber dichlorodiphenyltrichloroethane (DDT) and its metabolites (DDX) in fasting animals, but declined with DDX when animals were feeding. Our data show that POP burdens are high enough in seal pups to alter adipose function early in life, when fat deposition and mobilisation are vital. Such POP-induced alterations to adipose glucose use may significantly alter energy balance regulation in marine top predators with the potential for long term impacts on fitness and survival.

TOC/Abstract Art



Introduction

The impact of environmental and organismal accumulation of man-made chemicals is a growing global challenge^{1,2}. Persistent organic pollutants (POPs) are toxic³, accumulate in living tissues⁴, and resist biological and chemical degradation⁵. Widespread bans on their production initially lowered environmental POP levels⁶⁻⁹, but reductions have slowed or ceased more recently¹⁰⁻¹³. Legacy POPs are thus ubiquitous in the marine environment and remain a serious concern for top predators, in which biomagnification produces the highest burdens^{4,11,14-16}.

At high exposure levels POPs, such as polychlorinated biphenyls (PCBs), polybrominated diphenyl ethers (PBDEs) and organochlorine pesticides (OCPs), including dichlorodiphenyltrichloroethane (DDT), impair fertility, reproductive function, skeletal and neurological development, immunity and thyroid function in humans and wildlife¹⁷⁻²⁴. Recently, several POPs have also been implicated as lipid disruptors in humans and rodents^{25,26}. In fish and zooplankton, POPs can increase fat deposition and lipogenic gene expression²⁷⁻³¹. Conversely, exposure to POPs such as dioxins, lindane and OCPs can cause rapid energy reserve depletion, termed 'wasting syndrome'³²⁻³⁵. These apparently conflicting effects of different POPs on a variety of energetic processes highlight the need for further investigation of pollutant mixtures, doses and exposure routes, and their physiological consequences for organisms in natural environments.

POP-induced alterations to energy balance may have significant individual and population level consequences because fatness is often positively associated with reproductive fitness and survival³⁶. This is particularly true in marine mammals³⁷⁻³⁹, which rely heavily on subcutaneous fat reserves for insulation⁴⁰⁻⁴¹ and as a metabolic fuel when food is scarce. Many marine mammal species also undergo natural periods of

fasting during moulting, breeding, migration, or the postweaning fast that occurs in many phocid pups^{37,42-46}.

Marine mammals typically experience exceptionally high POP burdens due to their high fat content and top trophic level, and are regarded as sentinels of marine pollution^{14,16}. POPs are transferred to suckling young as they undergo rapid fat accretion⁴⁷. In fasting individuals, less hydrophobic POPs are released into the blood stream as lipid is mobilised and more recalcitrant POPs concentrate in the remaining blubber⁴⁷⁻⁵⁴. Phocid pups thus experience high POP exposure during both feeding and fasting, two fundamentally important periods for adipogenesis, fat mobilisation and maturation of energy balance pathways. Their reliance on adipose tissue to fuel metabolism during fasting or while searching for food in changing environments⁵⁵ place these top predators at high risk from adverse effects of POP exposure as a result of impaired fat cell function.

Marine mammals show structural changes in fatty acid (FA) content and thermal properties with blubber depth^{54,56-61}. Effects of blubber depth and nutritional state on metabolic function and sensitivity to POP exposure have not been explored. High POP levels are associated with altered adipose gene expression in seals⁶²⁻⁶⁴, polar bears (*Ursus maritimus*)⁵⁵, and killer whales (*Orca orcinus*)⁶⁵. POP cocktails from polar bear fat can induce adipogenesis in mouse (*Mus musculus*) pre-adipocyte cell lines and primary polar bear pre-adipocytes⁶⁶. However, the consequences of POP-induced gene expression changes for fat tissue function are unknown.

The major functions of fat tissue are to store lipid and release it for use as metabolic fuel. The balance between lipogenesis and lipolysis depends on the nutritional state of the organism and the hormonal milieu to which the tissue is exposed, and may be influenced by POPs. Glucose uptake, lactate production and glycerol release

are important metabolic properties of adipose tissue that contribute to whole animal glucose homeostasis and energy balance. Adipose tissue synthesises triacylglycerol (TAG) from glucose^{67,68}, such that glucose uptake rates can be used to investigate metabolic function of adipose tissue. Glucose uptake is altered by POPs at tissue and the whole animal level in other animals⁶⁹⁻⁸⁷. Of glucose taken up by adipose, 20% is stored as TAG, and 70% is used to synthesise lactate⁶⁷. Lactate production is thus an important route of glucose disposal and may be influenced by POP exposure. Glycerol is produced during lipolysis⁸⁸, and as a by-product of glycolysis⁶⁸, and is thus a major secretory product of adipose tissue. Unlike FA, glycerol cannot be re-esterified by adipose *in vitro*^{89,90}, making its production a useful marker of lipolytic⁸⁸, and, to some extent, glycolytic⁶⁸ activity.

We measured glucose uptake, lactate production and glycerol release as indices of metabolic activity in blubber by employing an explant approach we have developed using blubber biopsies from wild grey seals (*Halichoerus grypus*)⁹¹. Our approach allows experimental work to be performed *in vitro* in remote field locations to investigate fat tissue function⁹¹, which, in marine mammals, was previously limited to measurement of enzyme activities⁹². Here, we used blubber explants from wild grey seal pups, which remain on land during both feeding (during suckling) and fasting (during the post-weaning fast). We investigated whether metabolic properties of blubber differ by tissue depth, nutritional state and intrinsic POP burden. We also investigated acute changes in blubber metabolic characteristics in response to exogenous experimental treatments with high, but biologically relevant, POP concentrations.

Methods

Study site and animals

Field work was conducted on the Isle of May, Scotland (56°11'N, 02°33'W) from 7th November to 18th December 2016 under permit from Scottish Natural Heritage (SNH). Grey seal females that were part of a long-term study were observed from when they came ashore, and the date of birth of their pups was recorded. Pups were included if they were at an appropriate predicted mass for their age (number of days post-partum) late in the suckling period and if they did not have any obvious indications of ill health. We collected blubber biopsies for POP analysis and explant culture from 27 pups late in the suckling period(day 15). Pups were then left to wean naturally from their mothers, which occurred at approximately 18 days old. We then attempted to resample the same pups once during the postweaning fast (day 12 after weaning). As pups were still feeding for several days between the two sampling events, additional mass gain and increases in blubber lipid percentage were possible between the feeding/fasting sampling points. Four pups were removed from the study because they developed an unrelated infection. This gave a total of 27 pups (9 female and 18 male) sampled while feeding and 23 (7 female and 16 male) of the same animals sampled again during fasting.

Pups were defined as weaned based on daily observation to determine maternal departure date⁴³. Pups were then transferred to a large (~600m²) fenced off area of the island, away from the main colony, with access to a pool of fresh water, in accordance with UK Home Office guidelines and based on prior studies^{43,93}.

The sex of all pups was determined at early lactation (~ day 5 of suckling). Pups were weighed at early and late lactation; when they were initially transferred into the

pen (within 2 days after weaning); and every 5 days in the pen. Pups less than 30 kg at weaning were excluded from the study. To ensure pups were of a viable departure mass, early release criteria were set at loss of 25% of weaning mass or reaching 30kg, whichever was the higher mass⁴³. It was not necessary to release any of the study animals early. All pups were released into the colony with ready access to the sea.

All animal procedures were performed under the UK Home Office project licence #70/7806 and conformed to the UK Animals (Scientific Procedures) Act, 1986. All research received prior ethical approval from Abertay University and the University of St Andrews Animal Welfare and Ethics Committee.

Biopsy sampling

Biopsy sampling, transport, explant creation and culture protocols were adapted from a previous study⁹¹. Prior to sampling, pups were given a mass-specific intravenous dose of ZoletilTM and subcutaneous injections of LignolTM at the biopsy sites. Three full depth blubber biopsies, one 6 mm and two 10 mm, were taken⁹⁴. The 6 mm biopsy was immediately wrapped in foil and frozen at -20 °C for POP analysis^{47,50}. The two 10 mm biopsies were cut immediately, using sterile surgical scissors, into ‘inner’ (blubber closest to muscle layer) and ‘outer’ (blubber closest to the skin) tissue, and placed into separate 15 ml centrifuge tubes filled with warm (37 °C) sterile Krebs Ringer solution (pH 7.4; NaCl; 7.89 g/l; KCl; 0.373 g/l; MgSO₄; 0.12 g/l; K₂HPO₄; 0.07 g/l; Glucose; 0.99 g/l; HEPES; 4.77 g/l; CaCl₂; 0.11 g/l⁹⁵ with 1% Antibiotic Antimycotic Solution (all chemicals supplied by Sigma- Aldrich) for transport back to the field laboratory. A single hidden suture (Ethicon Vicryl Suture (W9114) Violet, 3-0, 20mm, 75cm with VICRYL Suture 1/2 circle ‘Taper Point Plus’ Needle) was used to close 10 mm biopsy sites, either vertically or horizontally through the skin⁹⁶. Pups were observed remotely

on a daily basis, and biopsy sites checked at each reweighing to document healing. The second sample was not taken until all the previous biopsy sites had healed.

Explant culture and POP exposure

All biopsies were returned to the field laboratory, processed and placed in the incubator within 30 minutes of collection, which is within timeframes used in prior explant studies on livestock^{97,98} and wild seals⁹¹. Upon arrival, tissue was washed with 1 ml sterile, warm Krebs Ringer buffer. Visible hair, muscle or blood contaminated tissue was removed. Tissue was minced using a sterile scalpel into 5-10 mg pieces, and 100 mg portions of tissue were weighed out in a sterile Petri dish. The 100 mg explants were dispensed into pre-prepared 12 well plates warmed to 37°C in a sterile, non-ducted PCR hood. In total, 1500µl of either complete cell culture media (medium 199, Hanks' Balanced Salts with 1% Antibiotic Antimycotic Solution, 1% FA supplement and 5% charcoal stripped foetal bovine serum) or complete media containing 150ng/ml POP mix (PCB Aroclor standard mix 1 (Sigma-Aldrich), containing Aroclor 1016; 500 µg/mL, Aroclor 1260; 500 µg/mL, Decachlorobiphenyl (PCB 209); 50 µg/mL and 2,4,5,6-Tetrachloro-m-xylene (an OCP); 50 µg/mL in acetone:methanol solution (2:3)) was added per well to generate control and +POP experiments. Plates were placed in a humidified incubator maintained at 37°C and 5% CO₂ (Thermo Scientific, Midi 40 CO₂ Incubator, model: 3404) for 24 h. All media was drawn off after 24 h and frozen at -20 °C. Explants were snap frozen in liquid nitrogen and stored at -80 °C.

Metabolite measurement

Glucose, lactate and glycerol concentrations were measured in media from explant experiments using Randox (County Antrim, UK) kits (glucose: GL364, lactate:

LC2389, glycerol: GY105) and standards in an RX Monza (Randox) Clinical Chemistry analyser (Model: 328-14-0914) as described previously⁹¹. Internal quality control measurements lay within $\pm 15\%$. Intra and inter assay variability for sample analysis was $< 5\%$. Mean variation between replicates from the same individual and of the same nutritional state and tissue type for the three metabolites analysed were as follows; glucose 26.6% (range 0.6-144%), lactate 46.9% (range 3-164%) and glycerol 19.8% (range 0.5-106%). Rates of glucose removal and accumulation of lactate (indices of glycolysis) and glycerol (indices of lipolysis) were calculated per 100 mg tissue⁻¹ hour⁻¹.

POP analysis

Extraction and detection of POPs took place at the Center for Analytical Research and Technology (CART), University of Liège, Belgium, using standard methods (supporting information SI 1). The 6 mm biopsies yielded 385 ± 7 (s.e.) mg blubber tissue per sample. Six non Dioxin-Like PCBs (NDL-PCBs) (28, 58, 101, 138, 153, 180), eight Dioxin-Like PCBs (DL-PCBs) (105, 114, 118, 123, 156, 157, 167, 189) and nine PBDEs (28, 47, 66, 85, 99, 100, 153, 154, 183) were analysed at CART in the Department of Chemistry. Four OCPs (o,p'-DDT, p,p'-DDT, dichlorodiphenyldichloroethane (DDD), dichlorodiphenyldichloroethylene (DDE)) were analysed at CART in the Laboratory of Animal Ecology and Ecotoxicology (LEAE).

POP concentrations were measured in ng g⁻¹ lipid, summed into Σ NDL-PCBs, Σ DL-PCBs, Σ PBDEs and Σ DDXs (o,p'-DDT, p,p'-DDT, DDD, DDE) and log transformed prior to statistical analysis.

Statistical analysis

Analyses were performed using the statistical package R 3.4.1⁹⁹. T-tests were used to compare log transformed POP concentrations between feeding and fasting samples. Pearson's correlation coefficient was calculated between all four Σ POP types (log transformed Σ NDL-PCBs, Σ DL-PCBs, Σ PBDEs and Σ DDXs). As all POP classes were highly significantly correlated, we explored the effects of each POP class separately. Generalised additive mixed models (GAMM)¹⁰⁰ were used to identify the variables that best explained glucose uptake, lactate production and glycerol production. Explanatory variables explored in these models included log transformed Σ NDL-PCBs, Σ DL-PCBs, Σ PBDEs and Σ DDXs in the blubber at time of sampling (each POP class tested separately in each model), the mass of the pup, the tissue depth (inner or outer), the nutritional state at sampling (feeding or fasting), the experimental conditions (control or +POP) and the pup's sex. Interaction terms between nutritional state, tissue type, experimental conditions and POP concentrations were also investigated (see SI Table 1). The model was fitted using the multiple generalized cross validation library *mgcv*¹⁰¹. The identities of individual pups were fitted as random effect smooths¹⁰² to account for the same individual generating multiple explants and any variation between replicates. The smoothing parameters were set by maximum likelihood to reduce the risk of overfitting¹⁰³. The models were fitted with a Gaussian error distribution. Glucose data were log transformed and glycerol data were square root transformed. No transformation was needed for lactate data. Model selection was performed by backwards stepwise elimination through examination of R^2 values, Akaike's information criterion (AIC) values, Δ AIC, Akaike's weights, QQ and residual plots

Results and Discussion

POP concentrations

Mean concentrations and ranges of the summed POPs in grey seal pup blubber are given in Table 1.

Table 1. Means and ranges for study pup masses and fat content of 6mm biopsies, with median concentrations and ranges of POPs in grey seal pup blubber in feeding and fasting states.

	Feeding mean	Feeding range	Fasting mean	Fasting range
Pup mass (kg)	41.9	31.4 – 52.6	40	32.8 – 50.4
Biopsy fat content (% lipid)	78.5	67.8 – 85.5	82.9	56.3 – 88.9
	Feeding median	Feeding range	Fasting median	Fasting range
Σ NDL-PCB (ng g ⁻¹ lipid)	354.29	194.2 – 790.5	541.73	302.5 – 1253.8
Σ DL-PCBs (ng g ⁻¹ lipid)	10.78	5.5 – 45.6	19.78	9.6 – 151.2
Σ PBDE (ng g ⁻¹ lipid)	13.81	6.4 – 57.9	20.2	10.2 – 70.1
Σ DDX (ng g ⁻¹ lipid)	169.09	98.5 – 420.8	209.56	128.4 – 345.7

Higher POP levels when pups were fasting (Welch's two sample t-test, $p < 0.001$ for all log transformed POP classes) are consistent with previous work showing that lipid mobilisation during fasting concentrates more lipophilic POPs in remaining fat tissue^{50,51,53}. Blubber may thus be more vulnerable to disruptive effects of contaminants during fasting if the POPs can interact with cellular machinery to exert negative effects.

Glucose uptake

Σ DL-PCBs, experimental conditions, tissue depth, and nutritional state best explained variation in glucose uptake, along with an interaction between tissue depth and nutritional state (GAMM: $R^2 = 0.36$, SI Table 2). Σ NDL-PCBs, Σ PBDEs and Σ DDX had no significant impact on glucose uptake (SI Table 1). Uptake was higher in inner compared to outer tissue ($p < 0.001$). Glucose uptake was lower during fasting than feeding in outer blubber ($p = 0.035$), but did not differ between nutritional states in inner tissue (Figure 1). In both inner and outer tissue, glucose uptake was inversely related to Σ DL-PCBs ($p < 0.001$) (Figure 1) and the slope of the relationship did not depend on nutritional state. Acute exogenous POP exposure increased glucose uptake compared to controls, (GAMM: $R^2 = 0.36$, $p < 0.001$, SI Table 2, Figure 2), and the increase was not affected by nutritional state or tissue depth. Doses of each PCB congener given in the explant POP exposure conditions, with the estimated percentage change for each PCB type measured in blubber are given in SI Table 3. A boxplot of glucose uptake data during different nutritional states and tissue depths is provided in SI Figure 1.




Figure 1. Scatter plot (left panel) and prediction line plot (right panel) showing the impact of intrinsic Σ DL-PCBs on glucose uptake in blubber from grey seal pups, comparing inner (black) /outer (blue) tissue and feeding (filled triangles or solid lines) /fasting (open circles or dashed lines) states (n = 27 feeding and n = 23 fasting pups). Individual was included as a random effect in the model.




Figure 2. Prediction line plots showing the impact of exposure to acute exogenous POPs on glucose uptake in blubber from grey seal pups, comparing inner (left panel) and

outer (right panel) tissue and feeding/fasting states across the range of intrinsic Σ DL-PCBs in the blubber. Individual was included as a random effect in the model.

These data show for the first time that metabolic activity in phocid blubber is altered by the POPs accumulated *in utero* and during suckling. POPs have been postulated to be ‘obesogens’ from epidemiological data in humans²⁴. Experimental studies have demonstrated that POPs can alter adipocyte function in rodents and human cell lines^{25,104}. POPs are associated with altered adipose and liver expression of metabolic pathway genes in adult polar bears^{55,66} and ringed seals (*Pusa hispida*)^{63,64}, but the functional consequences of such changes for fat tissue have not been identified. Recent experimental studies that replicate environmental or physiological POP concentrations and congener mixtures provide causal evidence of their immunological¹⁰⁵ and adipogenic effects⁹¹. Our work builds on these findings by showing that existing POP concentrations in blubber tissue, generated *in utero* and through lactational transfer, can alter metabolic properties of adipose tissue in young, developing seals.

Our data are consistent with experimental studies showing PCB 77, 153 and dioxin reduce glucose and insulin tolerance in mice, guinea pigs (*Cavia porcellus*), rabbits (*Oryctolagus cuniculus*) and in human cell lines⁶⁹⁻⁷⁷. PCBs are also associated with dysfunctional glucose homeostasis in humans⁷⁸⁻⁸¹. Here, only Σ DL-PCBs altered glucose uptake in blubber explants, consistent with an aryl hydrocarbon receptor (AhR)-dependent pathway in seals, as seen in other species^{71,78}. POP-induced reduced glucose use is accompanied by a fall in glucose transporter, GLUT4, mRNA expression and GLUT1 protein levels in other animals⁷⁵, and may also occur in seals. The opposite effects of an exogenous POP exposure to the intrinsic Σ DL-PCB levels on glucose

uptake may reflect the difference between short (overnight) *in vitro* and long-term (*in utero* and over 15 days of suckling) exposure, which may produce opposing effects on membrane fluidity and transport properties¹⁰⁶. Alternatively, either the DL-PCB dose in the acute exposure experiments may have been too low to affect glucose uptake (SI Table 3) or the exogenous POP cocktail produces different effects from the *in vivo* contaminant mixture. We cannot distinguish between these possibilities from our data.

Lower glucose uptake caused by DL-PCBs is likely to reduce fat deposition rate, which may be problematic for pups laying down fat while suckling. However, fat deposition may increase if acute exogenous POP exposure increases glucose uptake. Indeed, short-term exposure to a POP cocktail from polar bear blubber induced adipogenesis in a mouse pre-adipocyte cell line⁶⁶. The timecourse of POPs on both adipogenesis and lipogenesis needs to be explored.

NDL-PCBs, DDX and PBDEs in blubber did not affect glucose uptake, despite evidence of their impacts on glucose homeostasis in other animals^{82-87, 107}. Our data suggest that the negative relationship between PBDEs and first year survival of grey seal pups¹⁰⁴ is not driven by direct alterations to adipose metabolic properties. PBDEs may alter adipose function indirectly through endocrine disrupting effects, for example, by altering thyroid or insulin function^{86, 87, 108, 109}, highlighting the need for improved understanding of energy balance regulation in seals, and its disruption by POPs.

Glucose uptake was higher in inner tissue under all conditions, suggesting deeper blubber has greater metabolic activity than superficial tissue. This finding is consistent with regional stratification in structure; the theorised role of superficial blubber in thermoregulation; the possibility of smaller, more numerous adipocytes that are more metabolically active⁹² in deeper tissue; and the observation that deeper blubber

contains higher glycolytic enzyme levels compared to superficial blubber in fin whales (*Balaenoptera physalis*⁹²).

Glucose uptake fell as the animals transitioned from feeding to fasting in outer blubber, but not in inner blubber. This likely reflects high levels of fat deposition throughout the blubber during feeding, followed by cessation of fat accumulation during fasting. Metabolic rate falls in fasting pups^{42,110}, and this is likely reflected in reduced demands by all tissues, including blubber. Lower sensitivity to residual insulin present in stripped FBS may explain lower glucose uptake and greater decrease in outer tissue from fasting animals, a state in which many animals become less sensitive to the effect on insulin of glucose uptake¹¹¹⁻¹¹³. The lack of decreased glucose uptake in inner blubber in fasting animals is surprising. It is however plausible that cell membrane expression of GLUT4 and/or insulin -independent GLUT1, are reduced to a greater extent in outer compared to inner blubber during fasting. Indeed, fasting northern elephant seals (*Mirounga angustirostris*) maintain expression of GLUT4 in adipose cell membranes¹¹⁴.

Previous studies of metabolic effects of POPs on marine mammals have focussed on adults^{55,63,64,66}. Concentrations of POPs in adult seals are an order of magnitude greater than those found in pups, with adult seal blubber containing several thousand Σ PCBs (ng/g)¹¹⁵ compared to the levels detected in the pups here, which were in the hundreds of ng/g range (Table 1). Here, we have shown that pups have already accumulated high enough POP levels in their blubber to impact on adipose function at less than three weeks of age, when the ability to generate and regulate blubber is a key factor for survival¹¹⁶. Growing and dividing adipocytes are more vulnerable to disruption by DL-PCBs because they have higher AhR levels than mature, differentiated adipocytes^{117,118}. The sensitivity to POP-induced alterations to adipose

tissue metabolism is therefore likely to be greatest in young marine mammals, which are exposed to large quantities of POPs from their mothers^{47,50} and are likely to have a high proportion of proliferating pre-adipocytes as a result of rapid fat deposition¹¹⁹. The energy balance and survival consequences of reduced glucose uptake in young pups now needs to be explored.

Lactate production

Σ DL-PCBs, tissue depth, nutritional state, pup sex and interactions between Σ DL-PCBs and nutritional state, and tissue depth and nutritional state were retained in the model that best explained lactate production (SI Table 1 and 2). Σ NDL-PCBs, Σ PBDEs and Σ DDX had no significant impact on lactate production (SI Table 1). Lactate production increased with DL-PCBs in feeding pups, but decreased with DL-PCBs when pups were fasting (GAMM: $R^2 = 0.29$, $p = 0.003$; SI Table 2, Figure 3). Lactate generation was higher in males than in females ($p = 0.05$) and higher in inner tissue than outer tissue for any given DL-PCB load in any given nutritional state ($p = 0.001$) (Figure 3). The difference between inner and outer tissue tended to be greater during feeding than during fasting ($p = 0.055$). Lactate generation was not impacted by acute exogenous experimental POP exposure. A boxplot of lactate generation/uptake data during different nutritional states and tissue depths is provided in SI Figure 2.




Figure 3. Scatter plot (left panel) and prediction line plot (right panel) showing the impact of intrinsic Σ DL-PCBs on lactate generation/uptake in blubber from male grey seal pups (n=34), comparing inner (black) /outer (blue) tissue and feeding (filled triangles or solid lines) /fasting (open circles or dashed lines) states. Individual was included as a random effect in the model.

Lactate production is a major route of glucose disposal in fat tissue⁶⁷. However, the inhibitory effect of DL-PCBs on glucose uptake is only consistent with lactate production during fasting. During feeding, lactate production increased with increased DL-PCBs in the face of reduced glucose uptake. One explanation is that the electron transfer chain experiences increased push force during suckling, in combination with inhibition by DL-PCBs^{120,121}, which could elevate lactate production through increased reliance on glycolysis. Suckling pups with higher DL-PCB burdens could thus experience greater risk of metabolic acidosis from higher adipose lactate production, with attendant risk of adipose dysfunction, greater demand on liver gluconeogenesis by

to process the lactate, and thus heightened energetic costs that could impact on efficiency of mass transfer and fat accretion during suckling.

Adipose lactate production is a direct consequence of lactate dehydrogenase (LDH) activity¹²². LDH activity can fall during fasting¹²³, and LDH gene expression and enzyme activity also differ substantially between adipose depots in rats¹²². LDH activity can also be modulated by PCB exposure in rat and dog liver, and the effect is dose dependent^{124,125}. Shifts in LDH expression and activity may thus contribute to the differences in lactate production between depths and nutritional states seen here in response to DL-PCBs. Higher lactate generation in inner tissue and the greater difference between regions during feeding may be a result of greater glucose uptake and metabolic activity in inner blubber during suckling. Regional differences in metabolic properties may also relate to adipocyte size¹²⁶.

Higher lactate production by adipose from males than females has not been reported, either in adult seals⁹¹ or in humans and rodents¹²² and may result from the small number of females here. However, since adult females accumulate fat faster than adult males¹²⁷ and female juvenile northern elephant seals spare fat more effectively than males¹²⁸, it is possible that some metabolic differences are present in pups' blubber prior to overt size dimorphism. For example, females may use more lactate to synthesise FA¹²⁹ than males, releasing less net lactate.

Glycerol production

Nutritional state, an interaction between tissue depth and nutritional state, and an interaction between nutritional state and intrinsic blubber DDX were significant terms in the best model to explain glycerol production (SI Table 1 and 2, Figure 4). In addition, extrinsic POP treatment, mass and sex were retained as non significant

variables. Unexpectedly, glycerol production, was higher during feeding than fasting (GAMM: $R^2 = 0.74$, $p < 0.001$ Figure 4). In addition, lipolysis was higher in outer than in inner tissue during feeding, whereas inner tissue had a higher lipolytic rate than outer tissue in the fasted state ($p = 0.0003$; SI Table 2). Finally, lipolytic rate increased with intrinsic blubber DDX in fasting animals, whereas the opposite relationship between DDX and lipolytic rate occurred in feeding pups ($p < 0.0001$, Figure 4). A boxplot of glycerol generation data from different nutritional states and tissue depths is provided in SI Figure 3.




Figure 4. Scatter plot (left panel) and prediction line plot (right panel) showing the impact of intrinsic Σ DDX on glycerol generation in blubber from grey seal pups, comparing inner (black) /outer (blue) tissue and feeding (filled triangles or solid lines) /fasting (open circles or dashed lines) states ($n = 27$ feeding and $n = 23$ fasting pups). Individual was included as a random effect in the model.

POPs, such as dioxin and related compounds, and OCPs typically produce ‘wasting syndrome’ at high doses³²⁻³⁵, likely due to increased WAT lipolysis¹³⁰. In contrast, some studies on rat adipose or rodent cell lines show no effect of DDT or DDE

on lipolysis at lower doses^{131,132}. Our data show that lipolysis is increased in blubber of fasting grey seal pups in response to intrinsic tissue levels of DDT and its metabolites, but decreased in feeding pups. These data suggest, that, as in bats, seal pups in negative energy balance may be most vulnerable to the presence of DDX in their tissues³⁴. Interestingly, the increase in lipolysis is greater in outer tissue, which is proposed to be important for insulation⁵⁶⁻⁶¹, such that DDT may negatively affect both thermoregulatory and metabolic capability of blubber during fasting. A reduction in lipolytic rate with DDX in feeding animals could reflect inhibition of tissue turnover of FFA, increased lipogenesis or a shift towards lactate production rather than glycerol generation during glycolysis⁶⁸.

Lower glycerol production during fasting compared to feeding contrasts starkly with the typical fasting increase in lipolysis^{133,134}. Regional and nutritional differences may result from depth changes in metabolic activity and adipocyte size¹³⁵. Lower glycerol production in fasting pups could also arise from a reduction in aldolase activity¹³⁶. In addition, higher lipolytic rates in feeding animals may reflect their higher metabolic rate^{42,110}, faster rates of triglyceride turnover and/ or greater production of 3 carbon (3C) intermediates to protect tissues against high fuel availability and provide fuel for actively growing tissue elsewhere^{68,122}. High rates of glycerol production in feeding animals could protect adipocytes from metabolic acidosis, minimise overproduction of oxidising equivalents and maximise energy efficiency^{68,137}.

Value and wider application of the explant method.

Wildlife species are facing increasing anthropogenic threats that can limit their ability to find food and store fat as an energy reserve for lean times. Constrained, fragmented or reduced foraging ranges¹³⁸ and decreased prey availability^{139,140} limit

opportunities for fat accumulation, and force individuals to rely on adipose stores more frequently to sustain them until the next feeding opportunity. Understanding the impact of natural and anthropogenic stressors on energy balance requires experimental interventions that are practically and ethically challenging to accomplish in marine mammals and other wildlife species. *In vitro* methods are increasingly being explored as alternatives to whole animal work and lethal sampling. Prior efforts have generated viable skin tissue samples for culture via dermal punches and biopsy darts in both marine^{141,142} and terrestrial¹⁴³ wildlife species. Given the importance of fat tissue to the health and survival of wildlife in rapidly changing environments, there is a critical need for methods to better understand their adipose physiology. We previously used blubber explants from wild seals, which were returned to tissue culture facilities in a mainland laboratory⁹¹. Here, we collected and maintained viable blubber explants for 24 h in a remote field laboratory, with only core tissue culture equipment. The methodology outlined here could therefore be adapted for use in other wildlife species in remote field locations or where alternative *in vitro* methods are not available.

Outlook

Our explant method clearly shows promise in better understanding adipose tissue responses to contaminants and can be expanded beyond the parameters measured here. Although establishment of cell lines would permit a wider range of exposures, including time courses, dose responses and mixtures over longer duration experiments, such approaches have been hampered by the absence of typical proliferation and differentiation responses in primary cells from non-model species⁵³. A focus on understanding the requirements of such cell types and generation of cell lines would greatly facilitate progress in toxicological studies on adipose from wildlife species.

In summary, we have shown that dioxin-like PCBs disrupt adipose glucose uptake and lactate production, and DDX disrupt lipolysis in grey seal pups, using an experimental explant approach. Our results are particularly important because they demonstrate POP-induced alterations to adipose tissue function in young animals in the first weeks of life, for which rapid fat accumulation during suckling and mobilisation during fasting are vital for survival. They also highlight that different POPs may impact different metabolic characteristics, and that their effects are modified by nutritional state and tissue depth, producing complex effects on tissue function. Wider screening of metabolic properties of cells and tissue from non-model organisms is key to identify energy-balance disrupting effects of legacy and emerging contaminants. Our methods and findings also have wider application in understanding drivers of juvenile survival of other top marine predators, particularly in species that simultaneously experience high POP loads and rapid alterations in prey availability and habitat.

Acknowledgments:

We would like to thank Simon Moss, Holly Armstrong and everyone who assisted with colony observations and sample collection on the Isle of May during the 2016 season. We would like to thank Scottish National Heritage for the permit to work on the Isle of May and the skipper and crew of the Osprey and May Princess for help with gear and personnel transport. We would also like to thank Dr Georges Scholl, Catherine Adam and all of the staff at the University of Liège who provided training and assistance while KJR was in the lab processing samples for POP detection, and Dr James Turton at Abertay University for advice on adipocyte tissue cell culture techniques.

The long term program of research on grey seals at the Isle of May was funded by the SMRU national capability grant (SMRU 1001). KAB was funded by NERC grant NE/M013723/1 and AJH and KJR were funded by NE/M01357X/1 for this project. The funding bodies had no role in the design of the study, the collection of samples, the analysis of samples or data and the interpretation of the data. The authors declare that they have no competing financial interests.

Supporting Information

SI 1. Methodological details for POP detection.

SI Table 1. Candidate GAMM models exploring different summed POP types, potential interactions and the top models in the selection process.

SI Table 2. Model output from all final GAMMs analysing metabolic characteristics in explant experiments with their standard errors, estimates and p values. If predictor variables have parentheses, this indicates the tested group.

SI Table 3. Doses of each PCB type given in the explant PCB culture conditions, with the estimated percentage change for each PCB type measured in blubber tissue.

Σ DL-PCB within total dose: 1.6ng. Σ NDL-PCB within total dose: 202.6ng.

SI Figure 1. Glucose uptake in inner and outer blubber tissue from feeding and fasting grey seal pups (n = 27 feeding and n = 23 fasting pups) with median, upper and lower quartiles, 1.5 x interquartile range and outliers shown. Significant differences between

groups are detailed in the main text, as not all significant parameters and interactions influencing glucose uptake are plotted. Individual was included as a random effect in the model.

SI Figure 2. Lactate production in inner and outer blubber tissue from feeding and fasting grey seal pups (n = 27 feeding and n = 23 fasting pups) with median, upper and lower quartiles, 1.5x interquartile range and outliers shown. Significant differences between groups are detailed in the main text, as not all significant parameters and interactions influencing lactate generation/uptake are plotted. Individual was included as a random effect in the model.

SI Figure 3. Glycerol generation in inner and outer blubber tissue from feeding and fasting grey seal pups (n = 27 feeding and n = 23 fasting pups) with median, upper and lower quartiles, 1.5x interquartile range and outliers shown. Significant differences between groups are detailed in the main text, as not all significant parameters and interactions influencing glycerol generation are plotted. Individual was included as a random effect in the model.

References

1. Hop, H.; Borgå, K.; Gabrielsen, G. W.; Kleivane, L.; Skaare, J. U. Food web magnification of persistent organic pollutants in poikilotherms and homeotherms from the Barents Sea. *Environ. Sci. & Technol.* **2002**, *36*, 2589-2597.
2. Ashraf, M. A. Persistent organic pollutants; a global issue, a global challenge. *Env. Sci. Poll. Res.* **2017**, *24*, 4223-4227.

- 568 3. Van den Berg, M.; Birnbaum, L.S.; Denison, M.; De Vito, M.; Farland, W.;
569 Feeley, M.; Fiedler, H.; Hakansson, H.; Hanberg, A.; Haws, L.; Rose, M.
570 The 2005 World Health Organization reevaluation of human and mammalian
571 toxic equivalency factors for dioxins and dioxin-like compounds. *Toxicol.*
572 *Sci.* **2006**, *93*, 223-241.
- 573 4. Johnson-Restrepo, B.; Kannan, K.; Rapaport, D. P.; Rodan, B. D.
574 Polybrominated diphenyl ethers and polychlorinated biphenyls in human
575 adipose tissue from New York. *Environ. Sci. & Technol.* **2005**, *39*, 5177-
576 5182.
- 577 5. Jones, K. C.; De Voogt, P. Persistent organic pollutants (POPs): state of the
578 science. *Environ. Pollut.* **1999**, *100*, 209-221.
- 579 6. Fleet, R. R.; Plapp Jr, F. W. DDT residues in snakes decline since the DDT
580 ban. *Bull. Environ. Contam. Toxicol.* **1978**, *19*, 383-388.
- 581 7. ATSDR (Agency for Toxic Substances and Disease Registry). Toxicological
582 Profile for Polychlorinated Biphenyls. US Department of Health and Human
583 Services, Public Health Services: Atlanta, GA, **2000**;
584 <https://www.atsdr.cdc.gov/toxprofiles/tp.asp?id=142&tid=26#bookmark12>
- 585 8. Connell, D. W.; Miller, G.; Anderson, S. Chlorohydrocarbon pesticides in
586 the Australian marine environment after banning in the period from the
587 1970s to 1980s. *Mar Poll Bull.* **2002**, *45*, 78-83.
- 588 9. MacGregor, K.; Oliver, I. W.; Harris, L.; Ridgeway, I. M. Persistent organic
589 pollutants (PCB; DDT;HCH; HCB; and BDE) in eels (*Anguilla anguilla*) in
590 Scotland: current levels and temporal trends. *Environ. Pollut.* **2010**, *158*,
591 2402-2411.

- 592 10. French, T. D.; Campbell, L. M.; Jackson, D. A.; Casselman, J. M.; Scheider,
593 W. A.; Hayton, A. Long term changes in legacy trace organic contaminants
594 and mercury in Lake Ontario salmon in relation to source controls
595 trophodynamics and climatic variability. *Limnol Oceanogr.* **2006**, *51*, 2794-
596 2807.
- 597 11. Aguilar, A.; Borrell, A. DDT and PCB reduction in the western
598 Mediterranean from 1987 to 2002, as shown by levels in striped dolphins
599 (*Stenella coeruleoalba*). *Mar. Environ. Res.* **2005**, *59*, 391-404.
- 600 12. Mahmood, M.; Bhavsar, S. P.; Arhonditsis, G. B. Examination of temporal
601 trends in Lake Erie fish communities using dynamic linear modelling. *J*
602 *Great Lakes Res.* **2013**, *39*, 437-448.
- 603 13. Sutton, R.; Sedlak, M. D.; Yee, D.; Davis, J. A.; Crane, D.; Grace, R.;
604 Arsem, N. Declines in polybrominated di-ethyl ether contamination of San
605 Francisco Bay following production phase-outs and bans. *Environ. Sci.*
606 *Technol.* **2015**, *49*, 777-784.
- 607 14. Ross, P. S.; Ellis, G. M.; Ikonomou, M. G.; Barrett-Lennard, L.G.; Addison,
608 R.F. High PCB concentrations in free-ranging Pacific killer whales, *Orcinus*
609 *orca*: effects of age, sex and dietary preference. *Mar. Pollut. Bull.* **2000**, *40*,
610 504-515.
- 611 15. Goerke, H.; Weber, K.; Bornemann, H.; Ramdohr, S.; Plötz, J. Increasing
612 levels and biomagnification of persistent organic pollutants (POPs) in
613 Antarctic biota. *Mar. Pollut. Bull.* **2004**, *48*, 295-302.
- 614 16. Desforges, J. P.; Hall, A.; McConnell, B.; Rosing-Asvid, A.; Barber, J.L.;
615 Brownlow, A.; De Guise, S.; Eulaers, I.; Jepson, P.D.; Letcher, R.J.; Levin,

- 616 M. Predicting global killer whale population collapse from PCB pollution.
617 *Science*, **2018**, *361*, 1373-1376.
- 618 17. Damerud, P.O. Toxic effects of brominated flame retardants on man and
619 wildlife. *Environ. Internat.* **2003**, *29*, 841-843.
- 620 18. Toft, G.; Hagmar, L.; Giwercman, A.; Bonde, J.P Epidemiological evidence
621 of reproductive effects of persistent organochlorines in humans. *Reprod.*
622 *Toxicol.* **2004**, *19*, 5-26.
- 623 19. Fisk, A.T.; de Wit, C.A.; Wayland, M.; Kusk, Z.Z.; Burgess, N.; Robert,
624 R.; Braune, B.; Norstrom, R.; Blum, S.P.; Sandau, C.; Lie, E.; Larsen,
625 H.J.S.; Skaare, J.U.; Muir, D.C.G. An assessment of the toxicological
626 significance of anthropogenic contaminants in Canadian arctic wildlife. *Sci*
627 *Total Environ.* **2005**, *351*, 57-93.
- 628 20. Hammond, J. A.; Hall, A. J.; Dyrnka, E. A. Comparison of polychlorinated
629 biphenyl (PCB) induced effects on innate immune functions in harbour and
630 grey seals. *Aquat. Toxicol.* **2005**, *74*, 126-138.
- 631 21. Bonde, J. P.; Toft, G.; Rylander, L.; Rignell-Hydbom, A.; Giwercman, A.;
632 Spano, M.; Manicardi, G. C.; Bizzaro, D.; Ludwicki, J. K.; Zvezday, V.;
633 Bonefeld-Jorgensen, E. C.; Pederson, H. S.; Joensson, B. A. G.; Thulstrup,
634 A. M. Fertility and markers of male reproductive function in Inuit and
635 European populations spanning large contrasts in blood levels of persistent
636 organochlorines. *Environ Health Perspect.* **2008**, *116*, 269-277.
- 637 22. Chiu, A.; Chiu, N.; Beaubier, N. T.; Beaubier, J.; Nalesnik, R.; Singh, D.;
638 Hill, W. R.; Lau, C.; Riebow, J. Effects and mechanisms of PCB ecotoxicity
639 in food chains: Algae > fish > seal > polar bear. *J. Environ. Scie. Health C*
640 **2008**, *18*, 127-152.

- 641 23. Kramer, S.; Hikel, S. M.; Adams, K.; Hinds, D.; Moon, K. Current status of
642 epidemiological evidence linking polychlorinated biphenyls and non-
643 hodgkin lymphoma, and the role of immune dysregulation. *Environ. Health*
644 *Perspect.* **2012**, *120*, 1067- 1075.
- 645 24. Czerska, M.; Zieliński, M.; Kaminska, J.; Ligocka, D. Effects of
646 polybrominated diethyl ethers on thyroid hormones, neurodevelopment and
647 fertility in humans and rodents. *Int. J. Occupat. Med. Environ. Health* **2013**,
648 *26*, 498-510.
- 649 25. Grün, F.; Blumberg, B. Endocrine disrupters as obesogens. *Mol. Cell.*
650 *Endocrinol.* **2009**, *304*, 19-29.
- 651 26. Regnier, S. M.; Sargis, R. M. Adipocytes under assault: environmental
652 disruption of adipose physiology. *BBA-Mol. Basis Dis.* **2014**, *1842*, 520-533.
- 653 27. Yadetie, F.; Karlsen, O. A.; Eide, M.; Hogstrand, C.; Goksøyr, A. Liver
654 transcriptome analysis of Atlantic cod (*Gadus morhua*) exposed to PCB 153
655 indicates effects on cell cycle regulation and lipid metabolism. *BMC*
656 *Genomics* **2014**, *15*, 481.
- 657 28. Yadetie, F.; Oveland, E.; Doskeland, A.; Berven, F.; Goksøyr, A.; Karlsen,
658 O. A. Quantitative proteomics analysis reveals perturbation of lipid
659 metabolic pathways in the liver of Atlantic cod (*Gadus morhua*) treated with
660 PCB 153 *Aquat Toxicol.* **2017**, *185*, 19-28.
- 661 29. Speranza, E. D.; Colombo, M.; Tatone, L. M.; Cappelletti, N.; Mogoya,
662 M.C.; Colombo, J.C. Fatty acid alterations in the detritivorous *Prochilodus*
663 *lineatus* promoted by opportunistic feeding on sewage discharges in the Rio
664 de la Plata estuary. *J. Fish Biol.* **2016**, *89*, 2024-2037.

- 665 30. Lee, M. C.; Han, J.; Lee, S. H.; Kim, D. H.; Kang, H. M.; Won, E. J.;
666 Hwang, D. S.; Park, J. C.; Om, A. S.; Lee, J. S. A brominated flame
667 retardant 2, 2', 4, 4' tetrabrominated diphenyl ether (BDE-47) leads to
668 lipogenesis in the copepod *Tigriopus japonicus*. *Aquat Toxicol.* **2016**, *178*,
669 19-26.
- 670 31. Maisano, M.; Cappello, T.; Oliva, S.; Natalotto, A.; Gianetto, A.; Parrino,
671 V.; Battaglia, P.; Romeo, T.; Salvo, A.; Spano, N.; Mauceri, A. PCB and
672 OCP accumulation and evidence of hepatic alteration in the Atlantic bluefin
673 tuna, *T. thynnus*, from the Mediterranean Sea. *Mar Environ Res.* **2016**, *121*,
674 20-48.
- 675 32. Schwetz, B. A.; Norris, J. M.; Sparschu, G. L.; Rowe, V. K.; Gehring, P. J.;
676 Emerson, J. L.; Gerbig, C.G. Toxicology of chlorinated dibenzo-p-dioxins.
677 *Environ. Health Perspect.* **1973**, *5*, 87-99.
- 678 33. Clark Jr, D. R.; Prouty, R. M. Experimental feeding of DDE and PCB to
679 female big brown bats (*Eptesicus fuscus*). *J. Toxicol. Env. Health* **1977**, *2*,
680 917-928
- 681 34. Clark Jr, D. R.; Stafford, C. J. Effects of DDE and PCB (Aroclor 1260) on
682 experimentally poisoned female little brown bats (*Myotis Lucifugus*): Lethal
683 brain concentrations. *J. Toxicol. Env. Health* **1981**, *7*, 925-934.
- 684 35. Swanepoel, R. M.; Racey, P. A.; Shore, R. F.; Speakman, J. R. Energetic
685 effects of sublethal exposure to lindane on pipistrelle bats (*Pipistrellus*
686 *pipistrellus*). *Environ. Poll.* **1999**, *104*, 169-177.
- 687 36. Young, R.M. Fat, energy and mammalian survival. *Am. Zool.* **1976**, *16*, 699-
688 710.

37. Pomeroy, P. P.; Fedak, M. A.; Rothery, P.; Anderson, S. Consequences of maternal size for reproductive expenditure and pupping success of grey seals at North Rona, Scotland. *J. Appl. Ecol.* **1999**, *68*, 235–253.
38. Hall, A. J.; McConnell, B. J.; Barker, R. J. The effect of total immunoglobulin levels, mass and condition on the first-year survival of grey seal pups. *Funct. Ecol.* **2002**, *16*, 462–474.
39. Lidgard, D. C.; Boness, D. J.; Bowen, W. D.; McMillan, J. I. State dependent male mating tactics in the grey seal: the importance of body size. *Behav. Ecol.* **2005**, *16*, 541–549.
40. Irving, L.; Hart, J.S. The metabolism and insulation of seals as bare-skinned mammals in cold water. *Can. J. Zool.* **1957**, *35*, 497–511.
41. Rosen, D. A.; Renouf, D. Seasonal changes in blubber distribution in Atlantic harbor seals: indications of thermodynamic considerations. *Mar. Mammal Sci.* **1997**, *13*, 229–240.
42. Reilly, J. J. Adaptions to prolonged fasting in free-living weaned gray seal pups. *Am. J. Physiol. – Reg. I.* **1991**, *260*, 267–272.
43. Bennett, K. A.; Speakman, J. R.; Moss, S. E.; Pomeroy, P.; Fedak, M. A. Effects of mass and body composition on fasting fuel utilisation in grey seal pups (*Halichoerus grypus fabricius*): an experimental study using supplementary feeding. *J. Exp. Biol.* **2007**, *210*, 3043–3053.
44. Waugh, C. A.; Nichols, P. D.; Noad, M. C.; Nash, S. B. Lipid and fatty acid profiles of migrating Southern Hemisphere humpback whales *Megaptera novaeangliae*. *Mar. Ecol. Prog. Ser.* **2012**, *471*, 271–281.
45. Irvine, L. G.; Thums, M.; Hanson, C. E.; McMahon, C. R.; Hindell, M. A.. Quantifying the energy stores of capital breeding humpback whales and

714 income breeding sperm whales using historical whaling records. *Roy Soc*
715 *Open Sci.* **2017**, 4, 160290.

716 46. Oftedal, O. T. The adaptation of milk secretion to the constraints of fasting
717 in bears, seals, and baleen whales. *J. Dairy Sci.* **1993**, 76, 3234-3246.

718 47. Debier, C.; Pomeroy, P. P.; Dupont, C.; Joiris, C.; Comblin, V.; Le
719 Boulengé, E.; Larondelle, Y.; Thomé, J.P. Quantitative dynamics of PCB
720 transfer from mother to pup during lactation in UK grey seals *Halichoerus*
721 *grypus*. *Mar. Ecol. Prog. Ser.* **2003**, 247, 237-248.

722 48. Hall, A. J.; Gulland, F. M. D.; Ylitalo, G. M.; Greig, D. J.; Lowenstine, L.
723 Changes in blubber contaminant concentrations in California sea lions
724 (*Zalophus californianus*) associated with weight loss and gain during
725 rehabilitation. *Environ. Sci. Technol.* **2008**, 42, 4181-4187.

726 49. Tartu, S.; Bourgeon, S.; Aars, J.; Andersen, M.; Polder, A.; Thiemann, G.
727 W.; Welker, J. M.; Routti, H. Sea ice-associated decline in body condition
728 leads to increased concentrations of lipophilic pollutants in polar bears
729 (*Ursus maritimus*) from Svalbard, Norway. *Sci. Tot. Environ.* **2017**, 576,
730 409-419.

731 50. Debier, C.; Pomeroy, P. P.; Dupont, C.; Joiris, C.; Comblin, V.; Le
732 Boulengé, E.; Larondelle, Y.; Thomé, J.P. Dynamics of PCB transfer from
733 mother to pup during lactation in UK grey seals *Halichoerus grypus*:
734 differences in PCB profile between compartments of transfer and changes
735 during the lactation period. *Mar. Ecol. Prog. Ser.* **2003**, 247, 249-256.

736 51. Debier, C.; Chalon, C.; Le Boeuf, B. J.; De Tillesse, T.; Larondelle, J. P.
737 Mobilization of PCBs from blubber to blood in northern elephant seals

- 738 (Mirounga angustirostris) during the post-weaning fast. *Aquat Toxicol.*
 739 **2006**, 80, 149-157.
- 740 52. Peterson, S. H.; Hassrick, J. L.; Lafontaine, A.; Thomé, J. P.; Crocker, D. E.;
 741 Costa, D. P. Effects of age, adipose percent, and reproduction on PCB
 742 concentrations and profiles in an extreme fasting North Pacific marine
 743 mammal. *PLoS ONE* **2014**, 9, e96191.
- 744 53. Louis, C.; Tift, M.; Crocker, D. E.; Alexander, D.; Smith, D. R.; Debier, C.
 745 Isolation of progenitor cells from the blubber of northern elephant seals
 746 (*Mirounga angustirostris*) in order to obtain an *in vitro* adipocyte model—
 747 preliminary results. *Mar. Mammal Sci.* **2014**, 31, 764-773.
- 748 54. Louis, C.; Covaci, A.; Crocker, D. E.; Debier, C. Lipophilicity of PCBs and
 749 fatty acids determines their mobilisation from blubber of weaned northern
 750 elephant seal pups. *Sci Tot. Env.* **2016**, 541: 599-602.
- 751 55. Tartu, S.; Lille-Langøy, R.; Størseth, T. R.; Bourgeon, S.; Brunsvik, A.;
 752 Aars, J.; Goksøyr, A.; Jenssen, B. M.; Polder, A.; Thiemann, G. W.; Torget,
 753 V. Multiple-stressor effects in an apex predator: combined influence of
 754 pollutants and sea ice decline on lipid metabolism in polar bears. *Sci. Rep.*
 755 **2017**, 7, 16487.
- 756 56. Aguilar, A.; Borrell, A. Patterns of lipid content and stratification in the
 757 blubber of fin whales (*Balaenoptera physalus*). *J. Mammal.* **1990**, 71, 544-
 758 554.
- 759 57. Koopman, H. N.; Pabst, D. A.; Mclellan, W. A.; Dillaman, R. M.; Read, A.
 760 J. Changes in blubber distribution and morphology associated with
 761 starvation in the harbor porpoise (*Phocoena phocoena*): evidence for

762 regional differences in blubber structure and function. *Physiol. Biochem.*
763 *Zool.* **2002**, 75, 498-512.

764 58. Best, N. J.; Bradshaw, C. J.; Hindell, M. A.; Nichols, P. D. Vertical
765 stratification of fatty acids in the blubber of southern elephant seals
766 (*Mirounga leonina*): implications for diet analysis. *Comp. Biochem. Phys. B.*
767 **2003**, 134, 253-263.

768 59. Koopman, H. N. Phylogenetic, ecological, and ontogenetic factors
769 influencing the biochemical structure of the blubber of odontocetes. *Mar.*
770 *Biol.* **2007**, 151, 277-291.

771 60. Strandberg, U.; Käkälä, A.; Lydersen, C.; Kovacs, K. M.; Grahl-Nielsen, O.;
772 Hyvärinen, H.; Käkälä, R. Stratification, composition, and function of
773 marine mammal blubber: the ecology of fatty acids in marine mammals.
774 *Physiol. Biochem. Zool.* **2008**, 81, 473-485.

775 61. Liwanag, H. E.; Berta, A.; Costa, D. P.; Budge, S. M.; Williams, T. M.
776 Morphological and thermal properties of mammalian insulation: the
777 evolutionary transition to blubber in pinnipeds. *Biol. J. Linn. Soc.* **2012**, 107,
778 774-787.

779 62. Brown, T. M.; Ross, P. S.; Reimer, K. J.; Veldhoen, N.; Dangerfield, N. J.;
780 Fisk, A. T.; Helbing C. PCB related effects thresholds as derived through
781 gene transcript profiles in locally contaminated ringed seals (*Pusa hispida*).
782 *Environ. Sci. Technol.* **2015**, 48, 12952–12961

783 63. Brown, T. M.; Hammond, S. A.; Behsaz, B.; Veldhoen, N.; Birol, I.;
784 Helbing, C. De novo assembly of the ringed seal (*Pusa hispida*) blubber
785 transcriptome: a tool that enables identification of molecular health
786 indicators associated with PCB exposure. *Aquat Toxicol.* **2017**, 185, 48-57.

- 787 64. Castelli, M. G.; Rusten, M.; Goksøyr, A.; Routti, H. mRNA expression of
788 genes regulating lipid metabolism in ringed seals (*Pusa hispida*) from
789 differently polluted areas. *Aquat. Toxicol.* **2014**, *146*, 239-246.
- 790 65. Buckman, A. H.; Veldhoen, N.; Ellis, G.; Ford, J. K. B.; Helbing, C. C.;
791 Ross, P. S.. PCB-associated changes in mRNA expression in killer whales
792 (*Orcinus orca*) from the NE Pacific Ocean. *Environ. Sci. Technol.* **2011**, *45*,
793 10194-10202.
- 794 66. Routti, H.; Lille-Langøy, R.; Berg, M. K.; Fink, T.; Harju, M.; Kristiansen,
795 K.; Rostkowski, P.; Rusten, M.; Sylte, I.; Øygarden, L.; Goksøyr, A.
796 Environmental chemicals modulate polar bear (*Ursus maritimus*) peroxisome
797 proliferator-activated receptor gamma (PPARG) and adipogenesis in vitro.
798 *Environ. Sci. Technol.* **2016**, *50*, 10708-10720.
- 799 67. Marin, P.; Rebuffe-Scrive, M.; Smith, U.; Bjorntorp, P. Glucose uptake in
800 human adipose tissue. *Metabolis.* **1987**, *36*, 1154–1160.
- 801 68. Rotondo, F.; Ho-Palma A. C.; Remesar X.; Fernández-López, J.A.; del Mar
802 Romeero, M; Alemany, M. Glycerol is synthesized and secreted by
803 adipocytes to dispose of excess glucose, via glycerogenesis and increased
804 acyl-glycerol turnover. *Sci. Rep.* **2017**, *7*, Article number: 8983
- 805 69. Baker, N. A.; Karounos, M.; English, V.; Fang, J.; Wei, Y.; Stromberg, A.;
806 Sunkara, M.; Morris, A.J.; Swanson, H.I.; Cassis, L.A. Coplanar
807 polychlorinated biphenyls impair glucose homeostasis in lean C57BL/6 mice
808 and mitigate beneficial effects of weight loss on glucose homeostasis in
809 obese mice. *Environ. Health Persp.* **2013**, *121*, 105.
- 810 70. Baker, N. A.; English, V.; Sunkara, M.; Morris, A. J.; Pearson, K. J.; Cassis,
811 L.A. Resveratrol protects against polychlorinated biphenyl-mediated

812 impairment of glucose homeostasis in adipocytes. *J. Nutr. Biochem.* **2013**,
813 24, 2168-2174.

814 71. Baker, N. A.; Shoemaker, R.; English, V.; Larian, N.; Sunkara, M.; Morris,
815 A. J.; Walker, M.; Yiannikouris, F.; Cassis, L. A. Effects of adipocyte aryl
816 hydrocarbon receptor deficiency on PCB-induced disruption of glucose
817 homeostasis in lean and obese mice. *Environ. Health Persp.* **2015**, 123, 944.

818 72. Enan, E.; Matsumura, F. 2,3,7,8-Tetrachlorodibenzo-p-dioxin (TCDD)-
819 induced changes in glucose transporting activity in guinea pigs, mice and
820 rats in vivo and in vitro, *J. Biochem. Toxicol.* **1994**, 9, 97–106.

821 73. Enan, E.; Lasley, B.; Stewart, D.; Overstreet, J.; Vandevoot, C.A. 2,3,7,8-
822 Tetrachlorodibenzo-*p*-dioxin (TCDD) modulates function of human
823 luteinizing granulosa cells via cAMP signaling and early reduction of
824 glucose transporting activity. *Reprod. Toxicol.* **1996**, 10, 191–198.

825 74. Hsu, H. F.; Tsou, T. C.; Chao, H. R.; Kuo, Y. T.; Tsai, F. Y.; Yeh, S. V.
826 Effects of 2,3,7,8-tetrachlorodibenzo-p-dioxin on adipogenic differentiation
827 and insulin-induced glucose uptake in 3T3-L1 cells. *J Haz Mat.* **2010**, 182,
828 649-655.

829 75. Liu, P. C.; Matsumura, F. Differential effects of 2,3,7,8-tetrachlorodibenzo-
830 *p*-dioxin on the “adipose-type” and “brain-type” glucose transporters in
831 mice. *Mol. Pharmacol.* **1995**, 47, 65-73.

832 76. Kern, P. A.; Dicker-Brown, A.; Said, S. T.; Kennedy, R.; Fonseca, V. A. The
833 stimulation of tumor necrosis factor and inhibition of glucose transport and
834 lipoprotein lipase in adipose cells by 2,3,7,8-tetrachlorodibenzo-p-dioxin.
835 *Metabolis.* **2002**, 51, 65–68.

- 836 77. Wu, H.; Yu, W.; Meng, F.; Mi, J.; Peng, J.; Liu, J.; Zhang, X.; Hai, C.;
837 Wang, X. Polychlorinated biphenyls 153 induces metabolic dysfunction
838 through activation of ROS/ NF- κ B signalling via downregulation of HNF1b.
839 *Redox Biol.* **2017**, *12*, 300-310.
- 840 78. Arsenescu, V.; Arsenescu, R. I.; King, V.; Swanson, H.; Cassis, L. A.
841 Polychlorinated biphenyl-77 induces adipocyte differentiation and
842 proinflammatory adipokines and promotes obesity and atherosclerosis.
843 *Environ. Health Persp.* **2008**, *116*, 761.
- 844 79. Uemura, H.; Arisawa, K.; Hiyoshi, M.; Satoh, H.; Sumiyoshi, Y.; Morinaga,
845 K.; Kodama, K.; Suzuki, T. I.; Nagai, M.; Suzuki, T. Associations of
846 environmental exposure to dioxins with prevalent diabetes among general
847 inhabitants in Japan. *Environ. Res.* **2008**, *108*, 63-68.
- 848 80. Uemura, H.; Arisawa, K.; Hiyoshi, M.; Kitayama, A.; Takami, H.;
849 Sawachika, F.; Dakeshita, S.; Nii, K.; Satoh, H.; Sumiyoshi, Y.; Morinaga,
850 K. Prevalence of metabolic syndrome associated with body burden levels of
851 dioxin and related compounds among Japan's general population. *Environ.*
852 *Health Persp.* **2009**, *117*, 568.
- 853 81. Everett, C. J.; Mainous III, A. G.; Frithsen, I. L.; Player, M. S.; Matheson, E.
854 M. Association of polychlorinated biphenyls with hypertension in the 1999–
855 2002 National Health and Nutrition Examination Survey. *Environ. Res.*
856 **2008**, *108*, 94-97.
- 857 82. Camón, L.; Solà, C.; Martínez, E.; Sanfeliu, C.; Rodríguez-Farré, E.
858 Cerebral glucose uptake in lindane-treated rats. *Toxicol.* **1988**, *49*, 381-387.
- 859 83. Rivera, S.; Sanfeliu, C.; Rodríguez-Farré, E. Changes in regional brain
860 2[¹⁴C]deoxyglucose uptake induced in postnatal developing rats by single

861 and repeated nonconvulsant doses of lindane. *Pesticide Biochem Physiol.*
862 **1992**, *43*, 241-252.

863 84. Mahmood, A.; Agarwal, N.; Sanyal, S.; Subrahmanyam, D. Effects of DDT
864 (Chlorophenotane) Administration on Glucose Uptake and Brush Border
865 Enzymes in Monkey Intestine. *Acta Pharmacol. Tox.* **1978**, *43*, 99–102.

866 85. Norberto, S.; Calhau, C.; Pestana, D.; Faria A. Effects of Environmental
867 Pollutants on MCF-7 Cells: A Metabolic Approach. *J. Cell. Biochem.* **2017**,
868 *118*, 366-375

869 86. Howell III, G. E.; Meek, E.; Kilic, J.; Mohns, M.; Mulligan, C.; Chambers, J.
870 E. Exposure to p,p'-dichlorodiphenyldichloroethylene (DDE) induces fasting
871 hyperglycemia without insulin resistance in male C57BL/6H mice. *Toxicol.*
872 **2014**, *320*, 6-14.

873 87. Ruzzin, J.; Petersen, R.; Meugnier, E.; Madsen, L.; Lock, E.J.; Lillefosse,
874 H.; Ma, T.; Pesenti, S.; Sonne, S. B.; Marstrand, T. T.; Malde, M. K.
875 Persistent organic pollutant exposure leads to insulin resistance syndrome.
876 *Environ. Health Persp.* **2010**, *118*, 465.

877 88. Langin, D. Control of fatty acid and glycerol release in adipose tissue
878 lipolysis. *C. R. Biol.* **2006**, *329*, 598–607.

879 89. Vaughan, M. The production and release of glycerol by adipose tissue
880 incubated *in vitro*. *J. Biol. Chem.* **1962**, *237*, 3354–3358.

881 90. Chakrabarty, K.; Bombeck, C. T.; Sigel, B.; Tauber, J. W.; Jeffay, H.
882 Glycerokinase activity in human adipose-tissue as related to obesity. *Int. J.*
883 *Obesity* **1984**, *8*, 609–622.

884 91. Bennett, K. A.; Robinson, K. J.; Moss, S. E.; Millward, S.; Hall, A. J. Using
885 blubber explants to investigate adipose function in grey seals: glycolytic,

886 lipolytic and gene expression responses to glucose and hydrocortisone. *Sci.*
887 *Rep.* **2017**, 7, 7731

888 92. Pond, C. M.; Mattacks, C. A. The distribution, cellular structure, and
889 metabolism of adipose tissue in the fin whale, *Balaenoptera physalus*. *Can.*
890 *J. Zool.* **1988**, 66, 534-537.

891 93. Robinson, K. J.; Twiss, S. D.; Hazon, N.; Moss, S.; Pomeroy, P. P. Positive
892 social behaviours are induced and retained after oxytocin manipulations
893 mimicking endogenous concentrations in a wild mammal. *Proc. R. Soc. B.*
894 **2017**, 284, 20170554.

895 94. Bennett, K. A.; Hughes, J.; Stamatas, S.; Brand, S.; Foster, N. L.; Moss, S.
896 E. W.; Pomeroy, P. P. Adiponectin and insulin in gray seals during suckling
897 and fasting: relationship with nutritional state and body mass during nursing
898 in mothers and pups. *Physiol. Biochem. Zool.* **2015**, 201588, 295-310.

899 95. Cold Spring Harbor Protocol. Krebs-Ringer Modified Buffer (KRB) **2014**,
900 doi:10.1101/pdb.rec076406

901 96. Chaise, L. L.; Paterson, W.; Laske, T. G.; Gallon, S. L.; McCafferty, D. J.;
902 Théry, M.; Ancel, A.; Gilbert, C. Implantation of subcutaneous heart rate
903 data loggers in southern elephant seals (*Mirounga leonina*). *Polar Biol.*
904 **2017**, 40, 2307-2312.

905 97. Etherton, T. D.; Evock, C. M. Stimulation of lipogenesis in bovine adipose
906 tissue by insulin and insulin-like growth factor. *J. Anim. Sci.* **1986**, 62, 357-
907 362.

908 98. May, S. G.; Savell, J. W.; Lunt, D. K.; Wilson, J. J.; Laurenz, J. C.; Smith, S.
909 B. Evidence for preadipocyte proliferation during culture of subcutaneous

910 and intramuscular adipose tissues from Angus and Wagyu crossbred steers.
 911 *J. Anim. Sci.* **1994**, 72, 3110-3117.

912 99. R Development Core Team. R: A language and environment for statistical
 913 computing. R Foundation for Statistics Computing, Vienna, Austria. **2012**,
 914 <http://www.R-project.org>.

915 100. Wood, S. *Generalized Additive Models: An introduction with R*.
 916 Chapman and Hall/CRC: London, U.K. 2006.

917 101. Wood, S. mgcv: Mixed GAM Computation Vehicle with
 918 GCV/AIC/REML smoothness estimation. **2012**, [http://cran.r-](http://cran.r-project.org/web/packages/mgcv/index.html)
 919 [project.org/web/packages/mgcv/index.html](http://cran.r-project.org/web/packages/mgcv/index.html)

920 102. Wood, S. Low-rank scale-invariant tensor product smooths for
 921 generalized additive mixed models. *Biometrics* **2006**, 62, 1025-1036

922 103. Wood, S. Fast stable restricted maximum likelihood and marginal
 923 likelihood estimation of semiparametric generalized linear models. *J. Roy.*
 924 *Stat. Soc. B.* **2011**, 73, 3-36.

925 104. Lee, D. H.; Porta, M.; Jacobs Jr, D. R.; Vandenberg, L. N. Chlorinated
 926 persistent organic pollutants, obesity, and type 2 diabetes. *Endocr. Rev.*
 927 **2014**, 35, 557-601.

928 105. Desforges, J. P.; Levin, M.; Jasperse, L.; De Guise, S.; Eulaers, I.;
 929 Letcher, R. J.; Acquarone, M.; Nordøy, E.; Folkow, L. P.; Hammer Jensen,
 930 T.; Grøndahl, C. Effects of polar bear and killer whale derived contaminant
 931 cocktails on marine mammal immunity. *Environ. Sci. Technol.* **2017**, 51,
 932 11431-11439.

- 933 106. Totland, C.; Nerdal, W.; Steinkopf, S. Effects and location of coplanar
934 and noncoplanar PCB in a lipid bilayer: a solid state NMR study. *Environ*
935 *Sci Technol.* **2016**, *50*, 8290-8295.
- 936 107. Lim, J. S.; Lee, D. H.; Jacobs, D. R. Association of brominated flame
937 retardants with diabetes and metabolic syndrome in the US population,
938 2003–2004. *Diabetes Care* **2008**, *31*, 1802-1807.
- 939 108. Hall, A. J.; Kalantzi, O. I.; Thomas, G. O. Polybrominated diphenyl
940 ethers (PBDEs) in grey seals during their first year of life—are they thyroid
941 hormone endocrine disrupters? *Environ. Poll.* **2003**, *126*, 29-37.
- 942 109. Hall, A. J.; Thomas, G. O.; McConnell, B. J. Exposure to persistent
943 organic pollutants and first-year survival probability in gray seal pups.
944 *Environ. Sci. Technol.* **2009**, *43*, 6364-6369.
- 945 110. Nordøy, E. S.; Ingebretsen, O. C.; Blix, A. S. Depressed metabolism and
946 low protein catabolism in fasting grey seal pups. *Acta Physiol.* **1990**, *139*,
947 361-369.
- 948 111. Bryant, N.; Govers, R.; James, D. Regulated transport of the glucose
949 transporter GLUT4. *Nat. Rev Mol. Cell. Biol.* **2002**, *3*, 267-277.
- 950 112. Fowler, M. A.; Champagne, C. D.; Houser, D. S.; Crocker, D. E.
951 Hormonal regulation of glucose clearance in lactating northern elephant
952 seals (*Mirounga angustirostris*). *J. Exp. Biol.* **2008**, *211*, 2943-2949.
- 953 113. Houser, D. S.; Champagne, C. D.; Crocker, D. E. A non-traditional
954 model of the metabolic syndrome: the adaptive significance of insulin
955 resistance in fasting-adapted seals. *Front. Endocrinol.* **2013**, *4*, 164.
- 956 114. Viscarra, J. A.; Champagne, C. D.; Crocker, D. E.; Ortiz, R. M. 5' AMP-
957 activated protein kinase activity is increased in adipose tissue of northern

- 958 elephant seal pups during prolonged fasting-induced insulin resistance. *J.*
959 *Endocrinol.* **2011**, *209*, 317-325.
- 960 115. Vanden Berghe, M.; Weijs, L.; Habran S.; Das, K.; Bugli, C.; Rees, J. F.;
961 Pomeroy, P.; Covaci, A.; Debier, C. Selective transfer of persistent organic
962 pollutants and their metabolites in grey seals during lactation. *Environment*
963 *International* **2012**, *46*, 6-15.
- 964 116. Hall, A. J.; McConnell, B.J.; Barker, R.J. Factors affecting first - year
965 survival in grey seals and their implications for life history strategy. *J.*
966 *Animal Ecol.* **2001**, *70*, 138-149.
- 967 117. Shimba, S.; Todoroki, K.; Aoyagi, T.; Tezuka, M. Depletion of aryl-
968 hydrocarbon receptor during adipose differentiation in 3T3-L1 cells.
969 *Biochem. Biophys. Res. Commun.* **1998**, *249*, 131-137.
- 970 118. Shimba, S.; Wada, T.; Tezuka, M. Arylhydrocarbon receptor (AhR) is
971 involved in negative regulation of adipose differentiation in 3T3-L1 cells:
972 AhR inhibits adipose differentiation independently of dioxin. *J. Cell. Sci.*
973 **2001**, *114*, 2809-2817.
- 974 119. Sethi, J. K.; Vidal-Puig, A. J. Adipose Tissue Development, Structure
975 and Function. In *Metabolic Basis of Obesity*; Ahima, R. Eds.; Springer, New
976 York, NY 2011.
- 977 120. Chesney, C. F.; Allen, J. R. Oxidative phosphorylation and respiration by
978 liver mitochondria from polychlorinated biphenyl-intoxicated rats. *Biochem*
979 *Pharmacol.* **1974**, *23*, 1577-1582.
- 980 121. Nishihara, Y. Effects of polychlorinated biphenyls (Kanechlor 400) on
981 isolated rat liver mitochondria. *Arch. Environ. Contam. Toxicol.* **1983**, *12*,
982 517-522.

- 983 122. Arriafan, S.; Agnelli, S.; Sabater, D.; Remesar, X.; Fernández-López, J.
984 A.; Alemany, M. Evidences of basal lactate production in the main white
985 adipose tissue sites of rats. Effects of sex and a cafeteria diet. *PLoS ONE*
986 **2015**, *10*, e0119572.
- 987 123. Kolataj, A.; Rysińska, J.; Flak, P. The influence of selection on reaction
988 to stress in mice. III. Influence of fasting, immobilization and exposure to
989 cold on lactate dehydrogenase and aldolase activity in the liver and kidney. *J*
990 *Animal Breed Genet.* **1995**, *112*, 224-233.
- 991 124. Rao, C. V.; Banerji, S. A. Effect of feeding polychlorinated biphenyl
992 (Aroclor 1260) on hepatic enzymes of rats. *Ind. J. Exp. Biol.* **1990**, *28*: 149-
993 151.
- 994 125. Sonne, C.; Leifsson, P. S.; Dietz, R.; Kirkegaard, M.; Jensen, A. L.;
995 Shahmiri, S.; Letcher, R. J. Greenland sledge dogs (*Canis familiaris*)
996 develop liver lesions when exposed to a chronic and dietary low dose of an
997 environmental organohalogen cocktail. *Env. Res.* **2007**, *106*, 72-80.
- 998 126. King, J. L.; DiGirolamo, M. Lactate production from glucose and
999 response to insulin in perfused adipocytes from mesenteric and epididymal
1000 regions of lean and obese rats. *Obesity Res.* **1998**, *6*, 69-75
- 1001 127. Beck, C. A.; Bowen, W. D.; Iverson, S. J. Sex differences in the seasonal
1002 patterns of energy storage and expenditure in a phocid seal. *J. Animal Ecol.*
1003 **2003**, *72*, 280-291.
- 1004 128. Kelso, E. J.; Champagne, C. D.; Tift, M. S.; Houser, D. S.; Crocker, D.
1005 E. Sex differences in fuel use and metabolism during development in
1006 juvenile northern elephant seals. *J. Exp. Biol.* **2012**, *215*, 2637-2645.
- 1007 129. Katz, J.; Wals, P.A. Lipogenesis from lactate in rat adipose tissue.
1008 *Biochim. Biophys. Acta* **1974**, *348*, 344-356.

- 1009 130. Roth, W. L.; Weber, L. W. D.; Stahl, B. U.; Rozman, K. A
1010 pharmacodynamic model of triglyceride transport and deposition during feed
1011 deprivation or following treatment with 2,3,7,8-tetrachlorodibenzo-p-dioxin
1012 (TCDD) in the rat. *Toxicol Appl. Pharmacol.* **1993**, *120*, 126-137.
- 1013 131. Rao, T. N.; Subrahmanyam, D. Effect of DDT on adipose tissue lipolysis
1014 in rats. *Toxicol.* **1981**, *20*, 229-235.
- 1015 132. Howell III, G. E.; Mangum, L. Exposure to bioaccumulative
1016 organochlorine compounds alters adipogenesis, fatty acid uptake, and
1017 adipokine production in NIH3T3-L1 cells. *Toxicol. in Vitro* **2011**, *25*, 394-
1018 402.
- 1019 133. Sethi, J. K.; Vidal-Puig, A. J. Thematic review series: adipocyte biology.
1020 Adipose tissue function and plasticity orchestrate nutritional adaptation. *J.*
1021 *Lipid Res.* **2007**, *48*, 1253-1262.
- 1022 134. Duncan, R. E.; Ahmadian, M.; Jaworski, K.; Sarkadi-Nagy, E.; Sul, H. S.
1023 Regulation of lipolysis in adipocytes. *Annu. Rev. Nutr.* **2007**, *27*, 79-101.
- 1024 135. De Koster, J.; Van dem Broeck, W.; Hulpio, L.; Claeys, E.; Van
1025 Eetvelde, M.; Hermans, K.; Hostens, M.; Fievez, V.; Opsomer, G. Influence
1026 of adipocyte size and adipose depot on the in vitro lipolytic activity and
1027 insulin sensitivity of adipose tissue in dairy cows at the end of the dry
1028 period. *J. Dairy Sci.* **2016**, *99*, 2319-2328.
- 1029 136. Orevi, M.; Gorin, E.; Shafrir, E. Adaptive changes in
1030 phosphofructokinase and aldolase in adipose tissue. *Eur. J. Biochem.* **1972**,
1031 *30*, 418-426.
- 1032 137. Del Mar Romero, M.; Sabater, D.; Fernández-López, J. A.; Ramesar, X.;
1033 Alemany, M.. Glycerol production from glucose and fructose by 3T3-L1

- 1034 cells: a mechanism of adipocyte defence from excess substrate. *PLoS ONE*
 1035 **2015**, *10*, e0139502.
- 1036 138. Regehr, E. V.; Lunn, N. J.; Amstrup, S. C.; Stirling, I. A. N. Effects of
 1037 earlier sea ice breakup on survival and population size of polar bears in
 1038 western Hudson Bay. *J. Wildlife Manage.* **2007**, *71*, 2673-2683.
- 1039 139. Fa, J. E.; Ryan, S. F.; Bell, D. J. Hunting vulnerability, ecological
 1040 characteristics and harvest rates of bushmeat species in afrotropical forests.
 1041 *Biol. Conserv.* **2005**, *121*, 167-176.
- 1042 140. Cheung, W. W.; Sarmiento, J. L.; Dunne, J.; Frölicher, T. L.; Lam, V.
 1043 W.; Palomares, M. D.; Watson, R.; Pauly, D. Shrinking of fishes exacerbates
 1044 impacts of global ocean changes on marine ecosystems. *Nat. Clim. Change*
 1045 **2013**, *3*, 254-258.
- 1046 141. Mathews, E. A.; Keller, S.; Weiner, D. B. A method to collect and
 1047 process skin biopsies for cell culture from free-ranging gray whales
 1048 (*Eschrichtius robustus*). *Mar. Mammal Sci.* **1988**, *4*, 1-12.
- 1049 142. Marsili, L.; Fossi, M. C.; Neri, G.; Casini, S.; Gardi, C.; Palmeri, S.;
 1050 Tarquini, E.; Panigada, S. Skin biopsies for cell cultures from Mediterranean
 1051 free-ranging cetaceans. *Mar. Environ. Res.* **2000**, *50*, 523-526.
- 1052 143. Tovar, H.; Navarrete, F.; Rodríguez, L.; Skewes, O.; Castro, F. O. Cold
 1053 storage of biopsies from wild endangered native Chilean species in field
 1054 conditions and subsequent isolation of primary culture cell lines. *In Vitro*
 1055 *Cell. Dev-An.* **2008**, *44*, 309-320.